



# ROMAN CHAMOMILE



# Roman Chamomile

TIC



ピークテーブル TIC

ピーク#	保持時間	面積	面積%	高さ	高さ%	A/H	化合物
1	6.587	159662	2.02	83948	2.26	1.90	Propanoic acid, 2-methyl-, 2-methylpropyl ester
2	6.951	462536	5.86	278618	7.50	1.66	.alpha.-Pinene
3	7.061	89006	1.13	40890	1.10	2.18	2-Propenoic acid, 2-methyl-, 2-methylpropyl ester
4	7.277	55889	0.71	32192	0.87	1.74	Camphene
5	7.808	50872	0.64	27796	0.75	1.83	.beta.-Pinene
6	8.242	90488	1.15	46969	1.26	1.93	Butanoic acid, 2-methyl-, 2-methylpropyl ester
7	8.423	37499	0.48	20458	0.55	1.83	Propanoic acid, 2-methyl-, 3-methylbutyl ester
8	8.479	164858	2.09	78274	2.11	2.11	Butanoic acid, 2-methylbutyl ester
9	8.955	187747	2.38	58815	1.58	3.19	2-Propenoic acid, 2-methyl-, pentyl ester
10	9.099	1181250	14.97	621538	16.73	1.90	3-Methyl-2-butenic acid, heptyl ester
11	9.344	755671	9.57	323474	8.71	2.34	2(5H)-Furanone, 5-(2-methyl-2-propenyl)-
12	10.030	59747	0.76	29158	0.78	2.05	Butanoic acid, 2-methyl-, 2-methylbutyl ester
13	10.237	158603	2.01	80058	2.15	1.98	Propanoic acid, 2-methyl-, hexyl ester
14	10.731	201519	2.55	116607	3.14	1.73	Furan, 2,3-dihydro-3-methyl-
15	10.771	629199	7.97	322225	8.67	1.95	Ethyl 1-methylcyclopropanecarboxylate
16	10.836	1482355	18.78	510564	13.74	2.90	2-Butenoic acid, 3-methyl-, pentyl ester
17	11.149	329232	4.17	164067	4.42	2.01	2(10)-Pinen-3-one, (+/-)-
18	11.386	31503	0.40	16872	0.45	1.87	Bicyclo[3.1.1]heptan-3-one, 2,6,6-trimethyl-
19	11.690	114221	1.45	49379	1.33	2.31	(1R)-(-)-Myrtenal
20	12.435	1621869	20.55	799953	21.53	2.03	3-Methyl-2-butenic acid, hexadecyl ester
21	15.921	28522	0.36	13709	0.37	2.08	1,6-Cyclodecadiene, 1-methyl-5-methylene-8-(1-methylethyl)-, [s-(E,E)]-
		7892248	100.00	3715564	100.00		

※1 設備提供:株式会社 久留米リサーチ・パーク

※2 使用機器:島津製作所製 GC-MS:QP2010 Plus

※3 分析条件:サンプルは有機溶媒で500倍に希釈後、直接導入法にて分析。

サンプル前処理...アセトンで500倍希釈、カラム...DB-5ms、

昇温条件...40°C(2min)-10°C/min-240°C(5min)、NISTライブラリにて同定。